

ELITE

GX

Welcome to Your MyBody GX Personal Report

EliteGX Personal Report

July 29, 2018

Congratulations! You are holding in your hands the codes to unlock insights about your body that, up until now, have never been available. The science of the human body only recently has evolved enough to allow scientists to identify and analyze a person's DNA. This program not only provides you with a roadmap of your specific genes, but also gives direction on how you can potentially optimize your fitness and performance with this knowledge.

As an active, athletic adult, you carve out the time (and often make sacrifices) to prioritize exercise in your life. Whether your main motivation is being strong and having enough endurance to enjoy an active life; you like to participate in recreational sports and hobbies like skiing, soccer, and hiking; and/or you have aspirations to qualify for the Boston marathon, compete in a CrossFit competition, or even cross the finish line of an Ironman triathlon, you do the hard work to train your body to perform.

The explosion of technology designed to monitor your activity and fitness has made it easier than ever to plan and track your workouts. The Internet is filled with training plans. There are literally dozens of apps you can download that tell you what to eat and how to train to achieve your goals. Heart rate monitors let you track your workout intensity, while GPS equipped motion sensors tell you how fast and far you've gone. You can even buy sensors that evaluate your resting heart rate and tell you when you need to rest and when you're ready to go hard. However, tracking alone doesn't help you choose the type of training that will unlock your highest personal potential or the food that fuels you best or if you're more likely to get injured following a certain regimen.

It's no secret that not every workout plan and nutritional approach will work for every person. What has been secret until now is how to figure out the ones that most closely match your personal make up and maximize your potential. This report will help you to better understand the factors that can affect how your body works to get the most from your exercise and training time and efforts.

This report will provide you with results in 4 key areas that can affect the way your body responds to training. It includes in-depth analysis of your genotype for certain key genes that are related to what type of athlete you are; your predicted training response, optimum strategies for fueling for activity and fat burning; and your recovery and risk for injury.

What is Genetic Testing?

Genetic testing utilizes a physical specimen from the body (saliva, blood, or other tissues) to reveal information about a person's chromosomes or their genes. In addition to identifying key genes, information is evaluated about areas on each gene that may differ between people. These areas are known as single nucleotide polymorphisms (SNPs). We use the term genotype to describe the outcome of your individual genetic tests.

Which Traits Were Analyzed?

To produce your results, this program looks at genes that are related to 4 major categories: Mental & Physical Foundation; Training Response; Fuel Utilization, and Recovery & Risk for Injury. Some of the results are directly related to your cardiovascular and strength foundation and “trainability”—what types of exercise suit your genotype best. Some results are related to how to optimize your training by way of fueling, recovery, and risk management. Other results are relevant because they can affect your motivation and behaviors that support your workouts and training.

How Are Your Results Determined?

We provide a genetic analysis that indicates which gene combinations you have in each category. You will receive a rating based on our calculated score for each trait in a category. Some categories have only one gene associated with that trait; other categories have several genes associated with that trait. Our calculated score reflects the potential combined influences from one or more genes.

We also provide personalized training, fueling, and recovery tips based on the potential implications of these results. In most cases, the outcomes for a genotype are a response to a specific fueling strategy or exercise prescription.

For example, in the case of cardiovascular exercise response, we review the body of literature, pulling the most well-conducted, relevant studies. One large study may follow participants who performed 50 minutes of cardiovascular training 3 to 4 days per week for 5 to 6 months. Participants may have differed in their response to this regimen based on their genetics. Some may have experienced greater fitness gains, while others experienced smaller gains and showed a decreased ability to perform at higher effort levels. If your result suggests a more unfavorable response, this doesn't mean that you cannot improve your fitness performing cardiovascular exercise. You may simply need a slightly different approach to get more favorable results. That's where many exercisers get stuck, attempting to discern just what the ideal approach may be. But we have evaluated your potential genetic response and provided suggestions on how to enhance it based on evidence-grounded research recommendations, as well as the experience of our medical team.

This program uses the best available research on which to base your results. We have established stringent criteria for studies that we use to help us evaluate the potential impact of your genotype for each gene tested. There are many studies that include genetic analyses, but for a variety of reasons, not all of them are reliable or valid. In determining how to process your genetic analysis, we do not accept just any research that has been performed on a gene. We use the largest and most scientifically valid genome-wide association studies, in addition to other high quality research, to calculate a score for the different genes or gene combinations for all genes tested. There is still much to learn in the field of genetic analysis. We are choosing the best available research upon which to base our analysis and recommendations.

Why Is Your Genotype Important?

Your genotype reveals the blueprint for your body. The ratings we provide you with reflect your genotypes for each gene or set of genes. This shows you your potential response, based on your genetic analysis, to different aspects of performance and training response to exercise, nutrition, and lifestyle behaviors (e.g., how you might be affected by different exercise modalities and fueling strategies). Keep in mind that if your results show the presence of certain genotypes and your result suggests that you will exhibit either an “enhanced” or “below average” response, this does not mean that the outcome associated with that genotype is definitely how your body will or does react.

Your phenotype is the physical manifestation, or expression, of your genotype. But your phenotype may be different than your genotype—not all the genetic variations seen in an analysis are manifested. That’s because how the genes that you have are expressed is largely affected by your lifestyle and other environmental factors. While your analysis might show that you have an increased or decreased potential for a certain training or performance trait, it does not mean that you will, in fact, express that trait.

This is very important to keep in mind because there is a tendency to view genotype results as a definitive diagnosis and to assume that you absolutely have certain traits, when this is not what a genetic analysis measures. The analysis only measures your odds for different outcomes, or the likelihood that your phenotype will express what your genotype predicts. Your results only suggest that there is a greater or lesser chance that you may exhibit certain traits or responses. The fields of nutrigenomics and exercise genomics are new, but growing, areas of research. Much still needs to be known to understand about genes and their interactions with each other, and the role in which other influences such as diet, exercise and the environment play in whether you will express a trait associated with a certain genotype.

That said, results from a genetic analysis may provide insights into how your body might perform optimally. If you have a certain genotype for a specific trait, knowing how it might affect you and adjusting your training, nutrition, and behaviors to maximize this information could make a difference in getting better results from your training and fueling. We provide personalized suggestions that may help you achieve the best results from your fitness and training efforts. Our team considers the results of your genetic analysis, along with an analysis of personal factors that you report, which may also influence your training response and body composition, as well as evidence-based guidelines that suggest the most effective strategies for optimum performance. All of this information combined is used to determine which training and diet strategies and lifestyle behavioral changes may be most helpful to reach your goals.

What You’ll Learn About You

On the following pages, you will see a summary of your results. You’ll learn what your genotypes suggest about your ability to make muscle in response to strength training, to boost your VO2 Max (a gold standard measure of physical fitness) in response to cardiovascular workouts, to burn fat, and to use carbs and protein. You also will gain insights into your intrinsic motivation to exercise; your sensitivity to caffeine; your ability to recover and minimize inflammation; your injury risk, and more. Your analyzed genotype results are followed by a detailed explanation and success strategy. Our medical team has evaluated your potential response and taken into account what evidence-based research recommendations on nutrition, training, and lifestyle behaviors suggest are the best approach for optimum performance to provide you with concrete success strategies. This guidance may give you that extra edge in finding the right plan that helps you maximize the results you get from all your hard work. While we can’t change our genes, we can change our behaviors to take advantage of what our genes say about our bodies.

REPORT SUMMARY



MENTAL AND PHYSICAL FOUNDATION

Intrinsic Motivation To Exercise	MORE LIKELY	BDNF
Power and Endurance Potential	HIGHER POWER	ACTN3, AGT, IL-6, NOS3, ACE, FTO, IGF1, GNB3, IL6-174
Grip Strength and Muscular Fitness	ABOVE AVERAGE	TGFA, POLD3, ERP27, HOXB3, GLIS1, PEX14, LRPPRC, MGMT, SYT1, HLA, GBF1, KANSL1, SLC8A1, ACTG1, DEC1, IGFS9B
Testosterone Levels	AVERAGE	SHBG



TRAINING RESPONSE

VO2 Max	BELOW AVERAGE	AMPD1, APOE
Exercise Heart Rate Response	SLIGHTLY ABOVE AVERAGE	CREB1
Exercise Stroke Volume	NORMAL	KIF5B
Body Composition Response To Strength Training	ENHANCED	NRXN3, GNPDA2, LRRN6C, PRKD1, GPRC5B, SLC39A8, FTO, FLJ35779, MAP2K5, QPCTL-GIPR, NEGR1, LRP1B, MTCH2, MTIF3, RPL27A, EC16B, FAIM2, FANCL, ETV5, TFAP2B



FUEL UTILIZATION

Protein Utilization	ENHANCED	FTO
Fat Utilization	LOW	PPARG, TCF7L2, APOA5, CRY2, MTNR1B, PPM1K
Carb Utilization	NORMAL	IRS1
Caffeine Metabolism	NORMAL	AHR, RP11-10017.3-001, ARID3B, CYP1A1



RECOVERY AND INJURY RISK


Injury Risk	ABOVE AVERAGE	SPTBN1, MEPE, SLC25A13, MBL2/DKK1, LRP5, C18orf19
-------------	---------------	---

MENTAL AND PHYSICAL FOUNDATION

INTRINSIC MOTIVATION TO EXERCISE

WHAT YOUR GENES SAY ABOUT YOU:

Our analysis indicates that your genetic profile exhibits characteristics that make you **MORE LIKELY** to be intrinsically motivated to train. That means you are more likely to derive pleasure from participating in your sport or activity without much need for external reward. That means you are more likely to consistently stick to a routine and spend more time engaged in your activity, which in turn can make it easier to reap other rewards and performance benefits.



Your genetic profile indicates that you are **MORE LIKELY** to have intrinsic motivation to exercise.

You will be more inclined to maintain an exercise and training routine without the need for external motivation or rewards. Be sure to build time into your schedule to fit the training you want to do.

RELATED GENES / SNPs

BDNF

The gene and associated SNP included in this category has been shown to have significant associations with a person's intrinsic motivation to train.

Athletes participate and compete in their sports for a variety of reasons and each of us has our own personal motivations. Athletes who are intrinsically motivated are inclined to participate in a sport for internal reasons. They run marathons because they love to run. They push themselves because they are driven to see how good they can be. They enjoy the process of training with the outcome being secondary. Those who are not intrinsically motivated tend to be extrinsically motivated, or participate for external reasons, such as winning competitions, impressing peers, or in some cases material rewards like trophies, medals, and even cash and scholarships.

Intrinsic motivation may be embedded in your genes. In one study, researchers

SUCCESS STRATEGIES

Athletes who are intrinsically oriented tend to face fewer ups and downs in their motivation. They also tend to be less stressed when they hit training obstacles and have less anxiety over "losing" or disappointing outcomes, like missing a podium position or failing to hit a set goal. Generally speaking, they exercise longer and more often, because they find activity itself rewarding.

Being intrinsically motivated to exercise makes getting regular physical activity easier, but it doesn't mean you're immune to boredom or falling into a rut. These strategies will help keep your routine fresh and rewarding.

Challenge yourself. Maximize your exercise enjoyment by challenging yourself with new fitness goals on a regular basis. If you regularly run 10Ks, aim for a half or even full marathon. If you're into CrossFit, compete in a local competition. New challenges can add fuel to your intrinsic motivation to keep moving and keep you from falling into a training rut.

MENTAL AND PHYSICAL FOUNDATION

INTRINSIC MOTIVATION TO EXERCISE

Pay it forward. Use your exercise motivation for greater good (and be even more motivated to train, especially during times when it's harder to get going like cold, wet weather seasons) by signing up for a run, bike ride, or triathlon that benefits a charity of your choice. There are also apps that will donate money to your favorite charity for every step you take.

collected DNA samples from a group of healthy adult men and women then observed the group while they performed a 30-minute treadmill workout. After the half hour session was up, the exercisers were told that they had completed the session and they could either begin a cool down or could keep going if they wanted. Those with at least one copy of the met allele for the val66met polymorphism were more than 2 ½ times likely to keep going than their peers with a val/val genotype.


Our analysis investigated which genotype for this gene was present in your DNA. Your rating of either **MORE LIKELY** or **LESS LIKELY** reflects whether your genotypes included those that carried a risk for being low in intrinsic motivation or for being likely to be high in intrinsic motivation. Knowing that you're genetically more or less inclined to be intrinsically motivated can help you establish strategies that may help ensure your success.

MENTAL AND PHYSICAL FOUNDATION

POWER / ENDURANCE POTENTIAL

WHAT YOUR GENES SAY ABOUT YOU:

Our analysis indicates that your genetic profile exhibits characteristics that make you likely to have **HIGHER POWER** response to strength and power training.. You are likely to enjoy significant power gains in response to strength training. Your genes may help you exceed in sports where power plays a pivotal role, such as sprinting; ball sports such as baseball, football, tennis, and golf; track racing, competitive lifting, and CrossFit types of activities. You can capitalize on your genotype by prioritizing strength and power workouts in your exercise regimen.



Your genetic profile indicates that you are likely to have **HIGHER POWER** response to strength training than other genotypes

That means you are more likely to improve your power in response to strength training and excel in sports that are power based.

Power related sports involve faster, more forceful bursts of activity for a shorter time. Typical power sports are mixed martial arts, hockey, football, baseball, and CrossFit.

Whether you tend to be power or endurance oriented largely depends on your muscle fiber composition. Generally speaking, muscle fibers can be broken into two types: type I (slow twitch) and type II (fast twitch). Type I use oxygen for fuel, fire slowly, provide continuous energy, and have high endurance. Type II rely on anaerobic metabolism for fuel, fire rapidly, and are quicker to fatigue. Fast twitch fibers can further be broken down into type IIa, which are intermediate or hybrid fast twitch fibers in that they use both aerobic and anaerobic metabolism almost equally and type IIb, which are purely anaerobic and produce the highest rate of contraction for quick, powerful bursts of activity. They also fatigue very quickly.

RELATED GENES / SNPs

ACTN3, AGT, IL-6, NOS3, ACE, FTO, IGF1, GNB3, IL6-174

The genes and associated SNPs included in this category have been shown to have significant associations with a person's endurance and/or power potential, or how likely you are to have a positive response to aerobic endurance and/or power training, which in turn may help determine the activities at which you will be most successful.

A meta-analysis of 35 articles published between 2008 and 2016 that analyzed the DNA of 19,852 people identified nine genetic variations that have significant associations with being a power athlete. Other research has found that a specific allele of the ACE gene is heavily represented in endurance athletes like elite long distance cyclists and is beneficial for endurance, rather than power-related sports.

Knowledge of your genetic makeup can help you hone your training for the

MENTAL AND PHYSICAL FOUNDATION

POWER / ENDURANCE POTENTIAL

Literature shows that healthy, sedentary people tend to have a 50/50 split of type I and type II fibers. Power athletes are estimated to have up to 80 percent type II fibers. Likewise, elite marathoners may have 80 percent type I fibers. Your DNA largely determines your personal combination. While you can train and make adaptations to muscle fiber size and shape and, to a small extent type, research shows genetic predisposition accounts for greater than 60 percent with only about a third being influenced by training and nutrition.

SUCCESS STRATEGIES

Your genetic inclination to be positively responsive to power training gives you an advantage for using training to get strong and fast for power-oriented sports and activities. That also means, of course, that you're less likely to excel in endurance-based sports like distance running, long distance cycling, and 70.3 or Ironman level triathlon. Which isn't to say you won't be successful in those endeavors, but may be less likely than other endurance-oriented genotypes to compete at an elite level.

Do some heavy lifting. Research indicates that athletes with your genotype see optimal gains in response to high intensity, low repetition type strength training. Try performing multiple sets of low weight such as 5 sets of 3 to 4 reps of close to your max weight with generous amounts of recovery (about 4 minutes) between sets. When you're comfortable with that, shake it up and lift closer to your max by performing 10 sets of 2 repetitions. This type of lifting stimulates your neural drive, improves your intramuscular coordination (so the fibers in any given muscle work better in concert to generate force), stimulates growth hormone production, and generates myofibrillar hypertrophy, which means your muscle fibers become denser and stronger—all of which plays to the strengths of your genotype and will help you excel in power-based athletic endeavors.

Be explosive. The definition of power is work divided by time. The more work you can do in a short amount of time, the more powerful you are. You can think of it as taking your strength and applying speed, such as what you do when you sprint, swing a bat to knock a ball out of the park, and clean and jerk a heavy bar. To maximize your genetic tendency to build power, include explosive types of training like Olympic style lifts and CrossFit into your training. Explosive workouts train your fast-twitch muscles to fire powerfully and quickly.

optimum outcome. In a study published in *Biology of Sport*, researchers tested the power and endurance levels of 28 athletes from different sports and 39 soccer players. All the athletes underwent genetic testing and then were assigned to a training protocol that either matched their DNA analysis or did not match their DNA analysis. After 8 weeks, they retested the athletes' aerobic fitness and explosive power. Those who were in the DNA-matched training group performed significantly better than those who were not.

Our analysis investigated which genotype for these genes was present in your DNA. Your rating of **HIGHER ENDURANCE**, **EQUAL ENDURANCE/POWER**, or **HIGHER POWER**, reflects whether your genotypes included those that carried the likelihood of being more responsive to endurance training; equally responsive to endurance and power training, or more responsive to power training.

MENTAL AND PHYSICAL FOUNDATION

POWER / ENDURANCE POTENTIAL

HIIT it. It may be tempting to forgo cardiovascular exercise completely, and if you're training for power-based sports, you don't need much. (Obviously if you're looking to do a mud run or Spartan challenge type of event, you'll need to increase your cardio training to build endurance to complete the distance of the event). By practicing high intensity interval training (HIIT), you can take advantage of your genotype and get the best of both worlds. Intense intervals tap into and condition your type II, turbo fibers, which you have in abundance. Because your heart rate stays elevated you get cardiovascular fitness benefits.

In one head-to-head comparison, Canadian researchers found that cyclists who performed 30-minute sprint sessions three days a week that included six 30-second max efforts with 4 minutes rest reaped nearly identical fitness gains to a group who pedaled moderately 90 minutes to two hours three days a week. In a similar study, the same research team found that high-intensity sprint training also doubled time to exhaustion during a threshold test. Since your genotype is one that is more responsive to power-based exercise, HIIT is an efficient way to improve your cardiovascular fitness.

A good example of HIIT is tabatas. They're super short, but very demanding. You can do them while running, cycling, or on exercise equipment like an elliptical. To do them, warm up for 5 to 10 minutes. Then go as hard as possible (you're going for maximum power output) for 20 seconds. Recover at an easy pace for 10 seconds. Repeat 6 to 8 times. Rest 4 to 5 minutes. If you are accustomed to interval training repeat for another set or two. If you're new to intervals, stick to one set. Cool down for 5 to 10 minutes.


Rest and recover. Because your genotype rewards high intensity training it can be tempting to hit it hard every time you hit the gym or perform your activity of choice. Resist the urge. Remember that your body repairs and makes its metabolic adaptations when you rest and recover. Take a couple of days off from hard training each week. Support your training and recovery with a healthful diet, good lifestyle habits, and quality sleep. Consider incorporating yoga into your routine for cross training and recovery. It will help you maintain muscle and joint mobility, which improves performance and helps prevent injury.

MENTAL AND PHYSICAL FOUNDATION

GRIP STRENGTH / MUSCULAR FITNESS


WHAT YOUR GENES SAY ABOUT YOU:

Our analysis indicates that your genetic profile exhibits characteristics that make you likely to have **ABOVE AVERAGE** hand grip/intrinsic muscular strength. That means you are genetically likely to perform above average in tests of hand grip strength as well as tests of general muscular strength and endurance, such as sit ups, push ups and lower body exercises like squats and leg presses. That's good news because a strong grip is important for a range of sports like racquet and ball sports, strength training, and everyday activity, like carrying luggage, and overall muscular strength is a key component in many aspects of training and competing.



Your genetic profile indicates that you are likely to have **ABOVE AVERAGE** hand grip/intrinsic muscular strength.

You are genetically predisposed to perform above average on tests of grip strength as well as tests of general muscular strength and endurance.



RELATED GENES / SNPs

TGFA, POLD3, ERP27, HOXB3, GLIS1, PEX14, LRPPRC, MGMT, SYT1, HLA, GBF1, KANSL1, SLC8A1, ACTG1, DEC1, IGFS9B

The genes and associated SNPs included in this category have been shown to have significant associations with a person's grip strength, which in turn may help determine how successful you will be in activities requiring muscular strength and muscular endurance.

Hand-grip strength is not just about a firm handshake. It reveals a lot about your intrinsic muscular strength and fitness and may, when weak, also indicate an increased risk in fractures. So it's important not only for successful performance in many activities such as racquet and ball sports, resistance training, and off-road cycling, but also for general vitality and health.

Research has found that hand-grip strength is strongly correlated to muscular strength and endurance. In one study, significant correlations were found between grip

SUCCESS STRATEGIES

Being genetically inclined to have above average grip and intrinsic muscular strength does not guarantee you will be an elite level power or sprint athlete, but it may give you an advantage in sports where grip strength is a priority. It may also boost your ability to maximize muscular gains in the gym, since handgrip is often the weakest link when people try to lift heavy weights.

It's important to note, however, that everyone, regardless of genetic make up, has weaker hand grip strength than they used to. Your grip strength is partly determined by activity and lifestyle. People performing lots of manual labor will have stronger hands. Grip strength—along with the rise in technology and decline in manual labor—has also been in decline among younger Americans, according to research. One 2016 study of 237 men and women ages 20 to 34 published in the Journal of Hand Therapy found that men 25 to 29 years of age had grip strengths of 101 and 99 pounds of force (right and left hands

MENTAL AND PHYSICAL FOUNDATION

GRIP STRENGTH / MUSCULAR FITNESS

respectively) today, a loss of 26 and 19 pounds from 30 years ago. Young women lost roughly 10 pounds of force over the same time period.

Grip strength also naturally declines with age, so working to maintain your naturally stronger grip will help make training and tasks of daily living easier and to reduce the risk of falls in your older years, regardless of your genetic makeup.

Because you are genetically predisposed to have above average grip strength, you do not need to do special hand strengthening moves. However, you can maximize your natural ability by including moves that build grip and overall strength into your resistance training routine. One of the best is the farmer's carry move. To do it grab a pair of heavy kettlebells (or dumbbells if there are no kettlebells available) and grasp them firmly as you walk 30 feet, taking short quick strides. Put them down and rest. Repeat 3 times. Use the heaviest weight you can carry.

Other overall conditioning moves such as deadlifts, rows, Olympic style lifts like cleans and snatches, pull ups, and front squats will not only maximize your grip strength, but also total body strength, which gives you an advantage in any sport or activity.

strength and performance in tests of muscular strength and endurance including sit ups, push ups, leg extension, and leg press.

Grip strength is also highly heritable. A large-scale genome-wide association study including a combined sample of 195,180 men and women identified 16 SNPs associated with grip strength. A number of these are also associated with genes that are implicated in the structure and function of muscle fibers, which helps explain why grip strength is a good indicator of intrinsic overall muscular strength. The study also confirmed that these genetic determinants of muscle strength were linked to fracture risk, likely because low muscle strength increases risk of falling.


Our analysis investigated which genotype for these genes was present in your DNA. Your rating of **BELOW AVERAGE**, **NORMAL**, or **ABOVE AVERAGE** reflects whether your genotypes included those that carried the likelihood of having below average grip/intrinsic muscular strength, average intrinsic grip/muscular strength, or below average intrinsic grip/muscular strength.

MENTAL AND PHYSICAL FOUNDATION

TESTOSTERONE LEVELS

WHAT YOUR GENES SAY ABOUT YOU:

Our analysis indicates that your genetic profile exhibits characteristics that give you the **AVERAGE** likelihood of having low testosterone levels. Levels of the hormone naturally decline with age and low testosterone levels are not uncommon in even otherwise healthy men. According to the American Urological Association, about 4 out of 10 men over the age of 45 have low testosterone, otherwise known as low-T. The anabolic steroid hormone testosterone has been shown to improve muscle size, strength, and function and body composition and helps maintain overall quality of life and health. So it's important to try to maintain healthy levels throughout your lifespan



Your genetic profile indicates that you are at **AVERAGE** risk for having low testosterone levels.

So you are genetically predisposed to be at an average risk for low testosterone levels. Since testosterone levels normally decline with age, it is important to take actions to maintain healthy levels of this hormone.

Testosterone helps you maintain muscle size, strength, and function, as well as overall vitality and body composition and low testosterone is a common affliction, so even though your genotype doesn't put you at higher than normal risk for having low-T, it's important to maintain nutrition, exercise, and lifestyle habits that optimize your genotype and help you maintain healthy hormone levels throughout your life.

SUCCESS STRATEGIES

Being genetically inclined to be at average risk for low testosterone levels is good news in that you're not at a particularly high risk for low-T. However, since low testosterone levels are not uncommon in men over the age of 45 and testosterone is one of the metabolic keys to building lean muscle, burning fat, and maintaining good health and vitality, you should optimize your genotype by practicing hormone building and preserving nutrition, training, and lifestyle habits.

RELATED GENES / SNPs

SHBG

The genes and associated SNPs included in this category have been shown to have significant associations with testosterone levels in men, which in turn may impact your stamina, body composition, strength, mood, and ability to make and maintain lean muscle mass.

Testosterone is a steroid hormone that is secreted by the testes and adrenal glands. It is instrumental in determining muscle size, strength, and function and also plays a role in maintaining lower levels of body fat. Low testosterone levels (defined as less than 300 ng/dl) not only hinder your ability to make gains in the gym, but also can be harmful to your health, as it's been associated with heart disease, metabolic syndrome, type II diabetes, osteoporosis, muscle loss, and increased mortality risk.

Testosterone gradually declines after age 40 (a phenomenon sometimes referred to as "andropause").

MENTAL AND PHYSICAL FOUNDATION

TESTOSTERONE LEVELS

Maintain a healthy weight. As an active athlete, this is likely already a priority, but hormonal health is another reason to keep your weight in check, even during “off season” times when you may not be training as rigorously. The 2011 National Health and Nutrition Examination Survey of 1,265 men ages 20 to 90 reports that increases in BMI, waist circumference, and body fat were linked to relative decreases in testosterone levels. Research shows that men who are 20% over their ideal weight have 25 to 45 percent lower testosterone levels than their peers who have a healthy weight.

Perform compound moves. Strength training is the best way to increase testosterone levels and the more muscle you put in motion at one time, the bigger the boost in testosterone production. Favor compound moves like squats, kettlebell swings, and deadlifts, in your resistance training repertoire. Research shows this type of training is superior for firing up your body's testosterone production.

HIIT it. High volume cardio can increase stress levels and hinder testosterone output. If you don't already, include sprint-type, high intensity interval training (HIIT), which stimulates testosterone response, as part of your cardio workouts. Twice a week HIIT training may help your hormonal health and will make you stronger and faster to boot.

Respect your sleep. Your body goes into deep repair mode while you sleep, pumping out muscle mending and building anabolic hormones like human growth hormone and testosterone. Research shows that reduced sleep also can lead to low testosterone, as lack of sleep can deplete T-levels by up to 15 percent. Aim to get at least 6 hours a night, ideally closer to 7 to 8.

Know the signs. Low-T can cause a cluster of physical and psychological symptoms including anxiety, irritability, sleep disruptions, depression, memory impairment, fatigue, low libido, erectile dysfunction, and weight gain (especially around the waistline). If you notice one or more of these symptoms, talk with your health care provider.

Being overweight also lowers testosterone as does smoking, and excess alcohol consumption.

Testosterone levels are also largely hereditary. Studies in male twins indicate that genetic factors account for about 65 percent of the variation in serum testosterone. A recent genome-wide association study that included a combined sample of 14,429 men identified genotypes that were associated with serum testosterone levels. One specific genetic variation was associated with a 6.5 fold higher risk of having low serum testosterone, or a 30 percent prevalence of low testosterone in men with that genotype compared to only a 4.6 percent prevalence of low testosterone among men with a more favorable genotype for serum testosterone levels.


Our analysis investigated which genotype for these genes was present in your DNA. Your rating of **MORE LIKELY**, **AVERAGE**, or **LESS LIKELY**, or reflects whether your genotypes included those that carried the likelihood of having below average testosterone levels, average serum testosterone levels, or above average serum testosterone levels.

TRAINING RESPONSE

VO2 MAX

WHAT YOUR GENES SAY ABOUT YOU:

Our analysis indicates that your genetic profile exhibits a **BELOW AVERAGE** fitness response to high-intensity exercise. That means you may experience lesser gains in your VO2 Max from high intensity workouts as someone with a more favorable genotype. That doesn't mean you can't benefit from cardiovascular training, however. You can still improve your fitness and performance. But you will likely see greater gains from longer, moderate intensity workouts. You may also benefit from resistance training workouts that improve your muscular power, efficiency, and endurance.



Your genetic profile indicates that your fitness response to moderate-to-high-intensity cardio is **BELOW AVERAGE**.

Though your genotype might limit your absolute VO2 Max improvements in response to high-intensity exercise, you can still improve your cardiovascular endurance by performing longer, moderate intensity workouts and by including resistance training in your regimen.

Your muscles need oxygenated blood to generate fuel. The more work you ask them to do, the more fuel—and oxygen-rich blood—they need. As you start to exercise, your heart rate and your breathing increases and keeps increasing as the intensity rises, so you can send more and more oxygenated blood to your working muscles. Keep pushing and you'll reach a point where your body can't use any more—your system is tapped out. Your heart is pumping all it can and your muscles are using all they can. That is your VO2 Max, the scientific name for the measure of your oxygen capacity—how much oxygen-rich blood your heart can pump and how much your muscles can use per minute, which is expressed in ml/kg/min.

VO2 Max is influenced by your genes, but also by your size, gender and, because it naturally diminishes overtime, age. For example, a sedentary woman in her 30s or 40s may have a VO2 Max of about 26 ml/kg/min while an athletic woman of the same age will have VO2 Max closer to 56 ml/kg/min. A 50-year-old man in fair condition may have a VO2 Max of 30, while his cross country ski-racing friend has one of 55 ml/kg/min.

RELATED GENES / SNPs

AMPD1, APOE

The genes and associated SNPs included in this category have been shown to have significant associations with a person's cardiovascular fitness response to moderate-to-high intensity exercise.

VO2 Max is generally considered the best indicator of aerobic fitness and endurance potential. Factors that impact it are how strong and efficient your heart is, how well-developed your capillary system is to deliver blood into your muscles, and the size and number of the energy-producing furnaces known as mitochondria in your muscle cells. All of these factors—and in turn your VO2 Max—improve with moderate to high intensity training. People who are active will have a higher VO2 Max than their sedentary peers. It is also influenced by your size, gender and, because it naturally diminishes overtime, age.

How much you can improve your VO2 Max depends upon myriad factors, including

TRAINING RESPONSE

VO2 MAX

Some people appear more inclined to respond to higher intensity efforts while others respond to lower intensity efforts. In one study, researchers from Queen's University in Kingston, Ontario, and the University of Ottawa measured VO2 Max in a group of "non-responders." They then split the group in half and had one group perform classic endurance style moderate-intensity training for three weeks, while the other half performed Tabata-style, very high intensity interval training (HIIT) style workouts over the same period. After three weeks, they measured their VO2 Max levels, waited several months, and then brought the volunteers back to do another three-week training block, this time doing the opposite style training as before. In the end, about a third of the group had little response to endurance style training; a third had little response to HIIT training, but no one failed to respond at all and generally if they didn't respond to one type of training, they had a good response to the other style of training.

In a similar study, researchers had sedentary men and women begin a cardiovascular training routine that included up to 50 minutes of cardio machines, like spin bikes and treadmills, 3 to 4 days a week for 5 to 6 months. Those with an "unfavorable" fitness response to cardio genotype experienced smaller gains in their cardiovascular fitness from the training. However, they also were less able to perform high intensity efforts, suggesting that their optimal fitness response may be better achieved at a lower intensity of exercise.

You'll generally hit your VO2 Max upper limit within a year of consistent training. It's important to note that even among elite athletes, there is a wide variation in VO2 Max values and that just because you've hit your VO2 Max ceiling doesn't mean that you can't continue to improve your athletic performance. Raising your lactate threshold—being able to perform at a higher percentage of your VO2 Max before fatiguing—and improving your efficiency and economy (which allows you to save energy while cranking out a high pace) can help you make measurable gains regardless of absolute VO2 Max.

SUCCESS STRATEGIES

The good news is that even if your genotype doesn't respond optimally to a steady diet of high intensity workouts, you can still make gains and see results through an exercise routine that prioritizes longer endurance workouts. Skew your training to prioritize zone 2 and 3 workouts.

your current fitness level and the intensity of your training. Research finds that sedentary people who start training at about 75 percent of their max for at least 30 minutes 3 times a week can increase their VO2 Max an average of 15 to 20 percent over six months, but the range of response is large. Some people make enormous gains, while others make very few. The reason, we now know, is in your genes. We've also learned that, contrary to what was thought previously, there are very few actual exercise "non-responders." It's more a matter of to what type of cardiovascular intensity your body best responds.

Our analysis investigated which genotype for these genes was present in your DNA. Your rating of **NORMAL**, **BELOW AVERAGE** or **LOW** reflects whether your genotypes included those that carried a risk of reduced cardiovascular fitness response from moderate-to-higher-intensity exercise.

TRAINING RESPONSE

VO2 MAX

It's also important to realize that a high VO2 Max isn't necessarily a benchmark for success in longer endurance activities. Elite marathon runners tend to have lower VO2 Max values than short or middle-distance runners because the maximal amount of oxygen you can deliver becomes less important when you're performing for long durations at sub-maximum intensities. You may excel at longer endurance events and, of course, non-endurance events that don't require a high VO2 Max.


Also consider including resistance training to improve your muscular strength and economy, which can give you a large performance boost regardless of your VO2 Max. Research has found that endurance athletes like runners, cyclists, and triathletes who swap some cardiovascular training time for resistance training like CrossFit and plyometrics improve their performance in time trials and other cycling, running, and multisport events, despite having slightly lower total training hours.

TRAINING RESPONSE

EXERCISE HEART RATE RESPONSE

WHAT YOUR GENES SAY ABOUT YOU:

Our analysis indicates that your genetic profile exhibits characteristics that give you a likely **SLIGHTLY ABOVE AVERAGE** heart rate response to exercise. That means you are likely to experience a small to moderate decrease in your exercise heart rate with training. That does not limit your ability to make measurable fitness gains or mean that you will experience sub-par performance with training. Even small heart rate improvements are meaningful and heart rate is only one factor in exercise performance and success.



Your genetic profile indicates that you are likely to experience a **SLIGHTLY ABOVE AVERAGE** heart rate response to exercise.

You are likely to experience a small to moderate decrease in exercise heart rate with training. Though larger decreases may be advantageous, heart rate is only one measurement of fitness and performance potential.

RELATED GENES / SNPs

CREB1

The gene and associated SNP included in this category have been shown to have significant associations with a person's exercise heart rate response. Your heart's primary job is to keep your blood circulating, sending blood into the lungs to pick up fresh oxygen and then pumping out that oxygenated blood into the rest of the body so your cells can function. When you exercise, your heart pumps faster to keep your working muscles fueled.

As you become more fit, your body becomes more efficient at using oxygen so your heart rate doesn't have to rise as dramatically when you exercise. It also becomes lower when you're at rest. Having a lower resting heart rate doesn't only indicate better heart health, but also, because your heart can pump more oxygenated blood with fewer beats per minute, you have greater endurance and exercise capacity. Your genetics have a considerable influence over how

SUCCESS STRATEGIES

Athletes often prize a low resting, and subsequent lower exercising, heart rate as a sign of superior fitness. Though a decrease in beats per minute (bpm) is a sign of improved cardiovascular conditioning and a lower resting heart rate can be an indicator of good fitness, it is only part of a larger picture when it comes to performance. These strategies can help maximize your training with regards to your genetic inclination for a more moderate heart rate training response.

Know your numbers. Healthy adults have an average resting heart rate (RHR) of 60 to 80 bpm; RHR may be 100 bpm in sedentary adults and lower than 60 bpm for active adults. Because women are smaller, their average heart rate is up to 10 bpm higher.

TRAINING RESPONSE

EXERCISE HEART RATE RESPONSE

The first step is determining your current resting heart rate, because if you don't know where you're starting, you can't measure your progress. If you've been training for more than a few weeks, you may have already achieved a lower resting and exercising heart rate and will not see further declines. Keep in mind that research shows that if you naturally have a lower heart rate, you will not see as dramatic a decrease as someone who has a naturally higher heart rate might.

It's best to measure your resting heart rate first thing in the morning. You should be fully recovered from any recent hard training or racing, as that can elevate your morning heart rate. If you need to use the bathroom, do that first, so you're fully relaxed. Put on your heart rate monitor and measure for about a minute, noting your lowest heart rate number. If you don't have a heart rate monitor you can simply use a timer and place your fingertips on your pulse and count beats for a minute.

If you've just started training, check again in three to four weeks (again being sure you're fully recovered) to note any changes.

Consider your exercise intensities. Lower intensity exercise doesn't have as much of an impact on your resting and exercise heart rate as high intensity exercise. Research shows that one hour a week of high intensity aerobic training lowered resting heart rate more effectively than lower intensity bouts.

Practice other heart healthy behaviors. Exercise isn't the only thing that impacts your resting and exercising heart rate. Dehydration thickens your blood and raises your heart rate, so staying hydrated is key. Caffeine can make it higher, so sensible caffeine use will help keep a too high heart rate in check. Exercising in extreme temperatures will raise it. Your heart rate will be higher at high altitudes, especially if you're not acclimated.

Make it part of the larger picture. Resting heart rate is only one piece of your athletic portrait. Your training partner with a resting heart rate of 32 bpm may win a fitness contest on paper, but in the real world, you could still be stronger and faster and get to the finish line first. Training your strengths can help minimize any limitations.

dramatically your exercise heart rate responds to endurance training.

The HERITAGE Family Study of 472 men and women from 99 nuclear families found that after 20 weeks of endurance training, the average decrease in heart rate during steady state aerobic exercise (60% of VO₂ Max) was 11.3 beats per minute (bpm), but there was a large range among individuals, from a decrease of 42 bpm to an increase of 12 bpm. Variations in the CREB1 SNP were strongly associated with heart rate response to exercise, explaining about 20 percent of the variance in heart rate response.


Our analysis investigated which genotype for this gene was present in your DNA. Your rating of **NORMAL**, **SLIGHTLY ABOVE AVERAGE**, or **ABOVE AVERAGE** reflects whether your genotypes included those that make you more likely to have a small, small to moderate, or moderate decrease in exercise heart rate with training.

TRAINING RESPONSE

EXERCISE STROKE VOLUME

WHAT YOUR GENES SAY ABOUT YOU:

Our analysis indicates that your genetic profile exhibits characteristics that make you likely to have a **NORMAL** stroke volume response to exercise. That means you are likely to experience a typical increase in stroke volume in response to exercise training.



Your genetic profile indicates that you are likely to experience a **NORMAL** stroke volume response to exercise training.

A greater stroke volume response is advantageous as you can pump out more blood at a lower heart rate. However, stroke volume response is only one factor in exercise performance and success.

Untrained people have a stroke volume of about 50 to 70 ml/beat at rest, which increases to 110 to 130 ml/beat during high intensity efforts. Exercise makes your heart muscle bigger and stronger, so you have a greater stroke volume. The resting stroke volume in elite athletes averages 90 to 110 ml/beat (which is why their resting heart rate is also so low), which increases to as much as 150 to 220 ml/beat during high intensity exercise, according to research.

Your stroke volume response is also sport dependent. Swimmers generally see a smaller increase in stroke volume response than runners or cyclists; exercising heart rate is typically lower during swimming as well, because the supine position prevents blood from pooling in the lower extremities and there's less need for increased heart rate and stroke volume to meet the body's needs.

RELATED GENES / SNPs

KIF5B

The gene and associated SNP included in this category have been shown to have significant associations with a person's exercise stroke volume response.

There are two ways for your heart to get more oxygen-rich blood to your exercising muscles: pump faster (heart rate response) and pump out a greater volume of blood with every beat. The latter is your stroke volume response, the amount of blood ejected per beat from your left ventricle, as measured in ml/beat.

Stroke volume increases as your exercise intensity rises. How much your stroke volume improves with exercise is also largely hereditary. The HERITAGE Family Study of 483 men and women from 99 nuclear families found that after 20 weeks of endurance training, the average increase in stroke volume during steady state aerobic exercise (60% of

TRAINING RESPONSE

EXERCISE STROKE VOLUME

Increasing stroke volume is believed to be more efficient than increasing heart rate during exercise, as you can do more work at a lower heart rate with a higher stroke volume. That said, once you reach a certain intensity, your stroke volume plateaus and your heart rate increases to meet your increasing exercise demands.

Your genetic tendency to have an average stroke volume response to exercise training does not mean you cannot perform at a high level or successfully compete against someone with a larger stroke volume response, though you may need to work harder at a given heart rate.

You can minimize any disadvantages by training your sport-specific skills; developing muscular power and efficiency and other fitness elements not solely dependent on cardiovascular efficiency. It's also important that you maintain optimum hydration status, as dehydration diminishes blood volume, which can exacerbate the impact of a genetically lower stroke volume.

VO2 Max) was 3.9 ml/beat. But there was a large range of stroke volume response among individuals, ranging from a decrease of 41 ml/beat to an increase of 45 ml/beat. Variations in the KIF5B SNP were strongly associated with stroke volume response to exercise, explaining nearly 30 percent of the variance.


Our analysis investigated which genotype for this gene was present in your DNA. Your rating of **NORMAL** or **ABOVE AVERAGE** reflects whether your genotype included those that make you likely to have an average or above average stroke volume response to regular exercise training.

TRAINING RESPONSE

BODY COMPOSITION RESPONSE TO STRENGTH TRAINING

WHAT YOUR GENES SAY ABOUT YOU:

Our analysis indicates that your genetic profile exhibits an **ENHANCED** body composition response to resistance training exercise. That means that along with improving strength and building lean muscle tissue, you are likely to lose weight and lower your body fat when you engage in a regular strength training routine. That's good news because many sports favor high power to weight ratios—having more lean muscle and less fat generally raises that ratio and may give you an advantage. A healthy body composition also lowers your risk for chronic disease like heart disease and diabetes.



Your genetic profile indicates that your body composition response to strength training is **ENHANCED**.

That means you are more likely to both make muscle and lose fat when you strength training regularly. You can maximize the benefits of your favorable genotype by resistance training at least two to three times a week.

As the term suggests, body composition is what your body is made from, including bone, water, fat mass, and lean tissue. For purposes of training and athletic performance, we tend to focus on the amount of body fat you have in relation to muscle. For good health, men should strive for a body composition that is less than 25 percent fat and women should aim for less than 32 percent fat. Generally speaking, athletic men and women have lower body fat percentages with elite athletic males averaging 6 to 13 percent and women averaging 15 to 20 percent.

Different sports have widely different requirements in terms of ideal body composition. A football lineman who needs a lot of absolute mass will have a higher percentage of body fat (though still a huge quantity of muscle) than an elite level male triathlete, who will be slowed down by any extra weight that is not helping to generate power. Regardless of sport, maintaining a healthy body composition is

RELATED GENES / SNPs

XN3, GNPDA2, LRRN6C, PRKD1, GPRC5B, SLC39A8, FTO, FLJ35779, MAP2K5, QPCTL-GIPR, NEGR1, LRP1B, MTCH2, MTIF3, RPL27A, SEC16B, FAIM2, FANCL, ETV5, TFAP2B

The genes and their associated SNPs are included in this category have been shown to have significant associations with a person's ability to improve their body composition in response to strength training.

As you know, resistance training helps you build and maintain lean muscle tissue. It may also help reduce the percentage and sometimes amount of body fat you have. That overall improvement in body composition makes you stronger and quicker in most sports. A higher percent of muscle and lower percent of fat tissue also contributes to a leaner appearance and, potentially, to a higher metabolism, or greater number of calories burned each day.

TRAINING RESPONSE

BODY COMPOSITION RESPONSE TO STRENGTH TRAINING

advantageous as it can help lower your chances of developing cardiovascular disease as well as diabetes and certain cancers.

This genotype is particularly favorable for body builders and power-based athletes, but all athletes and active people benefit from a healthy body composition.

SUCCESS STRATEGIES

Strength training works best when you lift weight that is heavy enough to fully stimulate lean muscle tissue growth—something many people, including athletes—neglect to do.

Your genotype may assist in great body composition improvements by participating in a focused strength-based program like CrossFit or similar conditioning class. Otherwise, aim to perform full body strength training two to three days a week.

For the best results, continually challenge your muscles in novel ways, mixing up your sets and reps from session to session, so sometimes you're lifting heavy (3 to 4 sets of 3 to 6 reps, 3 to 5 minutes rest between sets); sometimes light (2 to 3 sets of 10 to 15 reps, 1 to 2 minutes rest between sets), and sometimes moderate (2 to 3 sets of 8 to 10 reps, 1 to 2 minutes rest between sets). The weight you choose should be heavy enough so the last reps in a set are very challenging (but you can still maintain good form). When the exercises become easy, add more weight to continue to obtain the benefits.

Numerous factors, including your predominant muscle fiber type (which you discovered in the first section), your hormones (including testosterone, also from the previous section in this report), and the type of strength training you do influence how your body composition will respond to a resistance training program. Your genotype also plays a significant role.

In one large study, researchers had 148 volunteers participate in an intense resistance-training program for one year. They found that those who carried the most "favorable" gene variations enjoyed a full gamut of body composition benefits and not only improved their strength and muscle mass, but also experienced significant weight loss and body fat reduction. Those with less favorable genotypes still got stronger, but showed a decreased ability to lose weight and reduce body fat percentage by resistance training.


Our analysis investigated which genotype for these genes was present in your DNA. Your rating of either **ENHANCED**, **NORMAL** or **BELOW AVERAGE** reflects whether your genotypes included those that carried a risk of an enhanced or reduced body composition response to resistance training exercise.

FUEL UTILIZATION

PROTEIN UTILIZATION

WHAT YOUR GENES SAY ABOUT YOU:

Our analysis indicates that your genetic profile exhibits an **ENHANCED** utilization of protein. Your score reflects the fact that your genotype does include the allele combination that resulted in greater weight loss when a higher percentage of protein was eaten. Studies that investigated this genotype found that a diet consisting of a higher percentage of protein resulted in optimal weight loss. This suggests that the amount of weight or body fat you lose when trying to get lean is very likely to be affected by the percentage of protein you eat.



Your genetic profile indicates that your response is **ENHANCED**.

You may be more likely to meet your weight loss and or maintenance goals by eating a moderate-to-high percentage of protein in your daily diet. Aim for 25% to 30% of your total calories to come from plant and or lean animal-based protein.

The good news is that people with this genotype lost weight on a moderate to high protein diet. However, people with this allele also lost more lean body mass compared to those without this genotype. If weight and body fat loss is a goal, be sure to lift weights if you don't already, to minimize or prevent the loss of muscle that can occur with weight loss.

SUCCESS STRATEGIES

To lose and or maintain weight, consume a diet that is moderate to high in protein and be sure to include strength training in your regimen to minimize the loss of lean tissue. Here's how to optimize your protein intake.

Skew higher if you're actively trying to lose weight. The body must get a certain minimum amount of protein for normal functioning, and that is considered to be around 10% of total daily calories when you are eating enough food to meet your daily energy needs. This minimum amount of protein must be eaten to support processes such as enzyme and hormone production, cell repair and synthesis of skin and hair cells. If you start

RELATED GENES / SNPs

FTO

The gene and associated SNP included in this category has consistently been shown to be associated with body fat mass and BMI. As an athlete, you need more protein than the average person, as protein is required for muscle growth and repair after training, workouts, and competing. Many active people also rely on a heavy protein intake to shed and maintain weight, with a large contingency adopting protein-centric eating plans like the Paleo diet in hopes to maximize lean body tissue and minimize fat. Low well that approach works may be largely dependent on your DNA.

One large study found that people with a specific FTO variant had more successful weight loss and shed more body fat, particularly high-risk abdominal fat, if they ate a moderate-to-high protein diet (25-30% of total daily calories) compared to a lower protein diet (15-20% of total daily calories), regardless of the

FUEL UTILIZATION

PROTEIN UTILIZATION

reducing your food intake to drop weight, you need to eat a slightly higher percentage of protein because you are eating less food overall. Your genotype suggests that, while losing weight, you may benefit from a higher percentage of protein – from 25% to 30%.

Maximize your essential amino acids. Protein in your foods should contain all of the essential amino acids, since your body requires these to produce proteins, as well as the other amino acids it uses to make compounds such as enzymes, hormones, and tissues in your body. Animal foods contain all of the essential amino acids in one food item, such as meat, poultry, fish or dairy products. But if your genetic analysis for the other macronutrients suggests that you should reduce your intake of total fat or saturated fat, choose leaner versions of animal foods or, better, opt for plant-based protein foods.

You can obtain all of the essential amino acids in many single plant foods, including grains such as quinoa, seeds, such as shelled hemp hearts (hemp seeds), and beans such as edamame or tofu. Or you can consume several complementary plant foods in the same day and obtain the essential amino acids your body needs (brown rice and black beans; nuts, grains and beans; veggies, beans and grains, etc.).

Adjust your intake to match your activity. For average people, the recommendation is to obtain between 0.8 and 1 gram of protein per 1 kilogram of body weight. So if you weigh 150 lbs., or 68 kg, it is recommended that you get between 54 and 68 grams of protein per day. You need more as an active person.

The Academy of Nutrition and Dietetics recommends that athletes who participate in light to moderate endurance training take in 1.2 to 1.7 grams of protein per kilogram, or about 0.55 to 0.8 grams of protein per pound of body weight each day. So that same 150-pound person would need 82 to 120 grams of protein a day.

Power and sprint athletes looking to gain muscle mass also need more than average amounts of protein. If you're one, the Academy recommends you take in 1.4 to 1.8 grams of protein per kilogram or about 0.64 to 0.82 grams of protein per pound of body weight daily to build muscle mass. That same 150-pound athlete who wants to build muscle should eat 96 to 123 grams of protein every day.

Finally, serious endurance athletes who perform high volume training that includes high intensity intervals require the most protein because of the rate they break down their muscle tissue. The Academy recommends that high intensity endurance

percentage of their that came from fat and carbohydrates. However, they also lost more non-fat mass which includes muscle with the weight loss, even though they were eating a higher protein diet and exercising.

Our analysis of your genes investigated which genotype for this SNP was present in your DNA. Your rating of either **NORMAL**, **SLIGHTLY ENHANCED**, or **ENHANCED** reflects whether your genotype included those alleles that exhibited protein sensitivity, because their presence suggests that you will be more successful in your weight loss and maintenance attempts by following a moderate to high protein diet.

FUEL UTILIZATION

PROTEIN UTILIZATION

athletes take in 1.4 to 2.0 grams of protein per kilogram, or about 0.7 to 0.9 grams of protein per pound of body weight every day. So that 150-pound, high-intensity endurance athlete may need to consume up to 135 grams of protein every day.

Spread it out. Eat a variety of high protein foods, including eggs, dairy products, poultry, meat, fish and seafood, legumes, soy products, nuts and seeds, and spread it out throughout the day. By eating protein with every meal, you ensure that your muscles always have what they need for repair and maintenance. Protein also improves satiety, so you're less likely to over eat.

It's a good idea to get a sense of how much protein you are getting by recording your food intake for at least a week and entering it into a diet app or online nutrition log that can calculate the percentage of each of the macronutrients that you eat. Then you can tweak your menu as needed to obtain your recommended percentage of protein.

Add heavy resistance training. Since this SNP is also associated with reduced lean body mass, such as muscle tissue, with weight loss, it is recommended that you include heavier weight training as part of your plan if weight loss is one of your goals. This may help minimize or prevent the loss of lean body mass that can occur with weight loss. Study your results for your genetic analysis for exercise-related genes for a more specific exercise prescription. But for optimal muscle strengthening, you should do exercises with weights targeting your major muscle groups. On 2 to 3, non-consecutive days per week, do 3 sets of 12 reps with weight heavy enough to feel "hard" or "very hard" by the end of each set.

FUEL UTILIZATION

FAT UTILIZATION


WHAT YOUR GENES SAY ABOUT YOU:

Our analysis indicates that your genetic profile exhibits a **LOW** utilization of fat. Your score reflects the fact that for the genes investigated, your genotype includes some of the unfavorable allele combinations. This means that you may be sensitive to the amount and type of fat in your diet. Research has shown that people with a similar genotype profile tend to have more body fat when they have more fat in their diet and they lose less weight when they are on a diet that contains a high amount of fat, especially saturated fat. This result also suggests that you may have a reduced level of fat oxidation, or fat-burning ability, when you eat a high fat diet. That's of concern for endurance athletes who depend on fat burning for optimum performance.



Your genetic profile indicates that your utilization of fat is **LOW**.

You may be sensitive to too much total fat and/or too much saturated fat in your diet. If getting lean is one of your athletic goals, you may experience less weight loss with a higher fat diet. Aim for a lower total and saturated fat diet.



RELATED GENES / SNPs

PPARG, TCF7L2, APOA5, CRY2, MTNR1B, PPM1K

The genes and their associated SNPs that are included in this category all have been shown in scientifically sound studies to have statistically significant associations with how sensitive people are to eating a diet high in fat.

Like most active people, you may be trying to optimize your body composition by losing body fat and increasing lean muscle. Some athletes, concerned about carbohydrate-related weight gain, will prioritize fat in their diets, sometimes opting for a diet that gets a high percentage of calories from fat. Studies show that genotype plays a large role in how the amount of fat in your diet affects weight loss.

One study found that those people with an unfavorable genotype were more likely to have more body fat, a larger waist size, and a higher BMI the more fat they ate, compared to others without the same genotypes.

SUCCESS STRATEGIES

Since your genes suggest that you may be sensitive to the fat in your diet and that you may be less efficient at burning fat when you eat a high fat diet, following a lower fat diet and keeping saturated fat to a minimum may help you to control your body weight and body fat, and to lose more weight if that is a goal.

So how much fat should you eat?

There are varying definitions of what is considered "low fat." Studies that look at dietary fat vary in how they quantify fat and there is no clear consensus on what constitutes a "high fat" vs. a "low fat" diet. The Acceptable Macronutrient Distribution Range (AMDR) for dietary fat that is recommended by the Institute of Medicine is a daily fat intake that is between 20% and 35% of total daily calories and it is recommended to eat less than 10% of calories from saturated fats.

FUEL UTILIZATION

FAT UTILIZATION

A “high fat” diet is usually considered to be one consisting of a percentage of fat intake on the upper end of the AMDR range, so from 30% to 40% of the day’s total calories. People who eat a lot of fast food and animal foods like meat and cheese can have fat intakes that are 50% or greater. However, some people who choose to eat a very low carb diet may consume up to 60% or 70% fat.

A “low fat” diet is usually considered to be one consisting of a percentage of fat intake that is on the lower end of the AMDR range, so from 15% to 25% of the day’s total calories. Since your genetic profile indicates that you might benefit from a lower-fat diet, it is suggested that you aim for the lower end of that fat intake range, keeping saturated fat to a minimum. Be sure to not go too low when cutting fat. Athletes need more fat than sedentary people not only to fuel activity, but also to assist in the production of essential steroid hormones, which control how your body responds to strenuous activity. Too little fat can cause hormone imbalances and hinder your athletic performance and recovery. Prioritize healthy, inflammation-reducing fats like omega-3 fatty acids found in fish, nuts and seeds, avocados, and olive oil over less healthful food sources like fried foods.

It’s tough to know how much fat you get unless you are actively tracking what you eat and entering it into a diet app or online nutrition log. You might find it helpful to first determine how much fat you are currently eating so that you can identify ways to decrease it to desired levels if it is too high. If you are eating more fat than is recommended, analyze what you eat and use the tips below to reduce the fat.

Easy ways to reduce your total fat

- Stick to a plant-based diet: Eat fewer — or cut out completely — animal foods (meat, poultry and dairy foods).
- If you eat animal foods, choose leaner or lower-fat versions. Since even lean meats still contain fat, including saturated fat, control portion sizes and avoid eating meat at every meal, or even every day.
- Substitute plant versions of animal foods: try almond, soy or coconut-based yogurts, substitute plant milks (soy, almond, rice, etc.) for dairy milk.
- Reduce the amount of oil you use, or omit it completely.

To reduce saturated fat

- Use healthy oils (sunflower, safflower, coconut) instead of butter or cream for cooking or seasoning.
- Choose plant-based spreads instead of using butter. Use peanut butter, hummus, pesto sauce, avocado, etc.

Another study found that people with a protective genotype appeared to be able to consume greater amounts of fat, but without exhibiting higher BMIs. Another study found that people who went on a higher fat, reduced calorie diet lost weight, but they lost less weight if they had an unfavorable genotype compared to those with a more favorable genotype.


Our analysis of your genes investigated which genotype for each of these 6 genes was present in your DNA. Your rating of either **NORMAL** or **LOW** reflects whether your genotypes included some or all of those that carried a risk of reduced weight loss ability from a diet that was high in fat.

FUEL UTILIZATION

CARB UTILIZATION

WHAT YOUR GENES SAY ABOUT YOU:

Our analysis indicates that your genetic profile exhibits a **NORMAL** utilization of complex carbohydrates. Your score reflects the fact that your genotype does not appear to produce greater weight loss with a higher complex carbohydrate diet, and you can expect to lose around the same amount of weight with either a low, moderate or higher complex carb diet. Weight maintenance and loss aside, you still need adequate carbohydrate intake to fuel the sports and activities you perform. Complex carbs provide the most nutrients, fiber and, long-lasting energy during exercise.



Your genetic profile indicates that your utilization of carbohydrates is **NORMAL**

A high carbohydrate diet may not be beneficial if your goal is to lose and/or maintain weight. You still need adequate carbohydrate intake to fuel your activities, however. Prioritize complex carbohydrates, which provide the most nutrients, fiber, and lasting energy.

As an athlete, you use stored glucose—glycogen—to fuel your activity, particularly high intensity activity. Your body relies on glucose for daily living, as well, and this is why blood sugar levels are maintained within a consistent range. In fact, brain cells and red blood cells use glucose as their primary source of energy. Cells also use fat as a fuel source, but to metabolize fat, there must be some glucose present to complete the process.

Glucose is a very important nutrient. But sometimes cells do not respond to the insulin being released, a condition known as insulin resistance. The result is the bloodstream can be overloaded with glucose. Chronic high blood glucose levels can lead to pre-diabetes and, if unchecked, eventually diabetes, or uncontrolled high blood sugar. People who are overweight and/or physically inactive are at higher risk of insulin resistance. Athletes, however, are not immune to insulin resistance: one study on amateur athletes found that 3 out of 10 had fasting blood glucose in the pre-diabetes range.

RELATED GENES / SNPs

IRS1

The gene and associated SNP included in this category has been shown to be associated with a person's insulin sensitivity and the potential effects of the amount of carbohydrates and fat in the diet. Insulin is a hormone released by the body that helps cells take in glucose, or sugar, for energy. Glucose is present in the blood after the digestion of carbohydrates from foods like fruits, vegetables, legumes and grains. Insulin is also released in response to eating protein as it helps to shuttle amino acids into cells.

The gene in this category seems to influence insulin resistance and the body's response to carbs in the diet. One long-term study found that people with a variant of this gene who ate a high carbohydrate, lower fat diet that consisted of high fiber, whole plant foods, as opposed to processed, lower fiber carbs, had greater insulin sensitivity—and lower levels of insulin and insulin resistance—and experienced greater

FUEL UTILIZATION

CARB UTILIZATION

Since carbohydrate intake triggers insulin release, many people, including athletes and recreationally active men and women, assume that eating more carbs is not healthy and can lead to body fat and weight gain, as well as diabetes. Athletes in sports like CrossFit include a large low-carb diet (such as Paleo) contingency. But the relationship is not that simple: many people who eat a high carbohydrate diet perform well, are not overweight, and do not have diabetes, and, in fact, may have much lower levels of blood glucose. Several large epidemiological studies have shown that increased carb intake actually leads to a lower risk of diabetes and that, surprisingly, increased protein intake, increases the diabetes risk.

The types of carbs you eat play a role: Complex carbohydrates or starches are those that are made up of sugar molecules that are strung together in long, complex chains, as opposed to simple carbs, which are simple sugars like fructose and glucose. Complex carbs take longer for your body to digest, so have less of a “spike” effect on blood glucose levels. Sources include: whole grains like brown rice, quinoa, oatmeal, barley, bulgur, and buckwheat, vegetables, fruits, and legumes. If you eat mostly processed carbs (as opposed to fiber-rich complex carbs), you are likely to release greater amounts of insulin and this could affect your insulin resistance.

weight loss compared to eating a lower carb, higher fat diet. This is especially important as an active person, as maintaining healthy insulin sensitivity allows you to store the carbohydrates you eat for energy as well as to promote recovery after exercise and training.

Our analysis of your genes investigated which genotype for this gene was present in your DNA. Your rating of either **NORMAL** or **ENHANCED** reflects whether your genotype included those genes that improved insulin sensitivity and weight loss from a higher carb and slightly lower fat diet.

SUCCESS STRATEGIES

For weight loss, your genotype suggests that a high carbohydrate diet is not necessarily beneficial. Though there are no universally accepted definitions, high carb diets generally include greater than 60 percent of total calories from carbohydrates, and low carb diets generally include less than 30 percent of total calories from carbohydrates. Review your results from the other macronutrient genetic traits for more guidance on the weight/fat loss diet composition that might work best for you.

Performance-wise, your genotype does not suggest improved insulin sensitivity with high-carb intake. That does not mean you should eschew carbohydrates all together, however. But rather follow the carbohydrate intake recommendations found in current American College of Sports Medicine position statement on nutrition and athletic performance: reserve your higher carbohydrate intake for hard training days, races, and competitions. On easy or rest days, lean toward low carbohydrate intake.


Eat the majority of your carbohydrates from complex carbohydrate sources, which are highest in fiber, nutrients, and are best for your general health. Save simple carbohydrates like refined bread, pasta, sweets, and gels for when you need quick bursts of energy, such as right before a race or during prolonged activity, when you need fuel that is quickly absorbed. During such activity, you can aim for about 30 to 60 grams (120 to 240 calories) of carbohydrates per hour after the first hour to 90 minutes of activity.

FUEL UTILIZATION

CAFFEINE METABOLISM

WHAT YOUR GENES SAY ABOUT YOU:

Our analysis indicates that your genetic profile exhibits a **NORMAL** rate of caffeine metabolism. Having this genotype means you can enjoy the ergogenic benefits of this stimulant that include increased alertness, improved fat burning and glycogen sparing, lower rate of perceived exertion during hard efforts, and increased time to fatigue, especially during endurance events.



Your genetic profile indicates that you are likely to have a **NORMAL** rate of caffeine metabolism.

Your liver breaks down and clears the stimulant at a normal rate, and you are likely to benefit from using caffeine as an ergogenic aid in the sports and activities of your choice.

RELATED GENES / SNPs

AHR, RP11-10017.3-001, ARID3B, CYP1A1

The genes and their associated SNPs that are included in this category have been shown to have significant associations with a person's ability to metabolize caffeine.

Caffeine is well known and widely used as a legal stimulant. On the endurance front, caffeine increases the body's ability to use stored fat as fuel, which spares limited muscle glycogen (stored carbohydrate) stores. It also increases beta-endorphins to enhance feelings of wellness while also lowering your perceived exertion, so hard efforts feel easier. However, not everyone responds equally...or favorably. Some people suffer from negative caffeine side effects after one ill-timed cup of coffee, while others can drink several cups a day and feel fine.

We now know this disparity is largely hereditary. Caffeine is rapidly absorbed into the bloodstream, with levels peaking

Research dating back to the '70s has consistently shown that caffeine can improve sports performance, particularly endurance performance, where the average improvement in exercise trials is about 24 percent in time to exhaustion and 3.1 percent in time to completion. It may also improve muscle power and endurance for power and sprint-based sports.

Caffeine primarily interacts with adenosine, a chemical in your central nervous system that regulates sleeping and waking. As adenosine accumulates, it inhibits nerve activity and causes drowsiness. Caffeine essentially blocks adenosine, preventing your nerve activity from slowing down, which increases alertness and brain activity and reduces tiredness, which benefits all sports performance. It also increases circulating epinephrine, the hormone responsible for your fight or flight response, which helps you feel physically and mentally keyed up to perform.

SUCCESS STRATEGIES

For maximum benefit, ingest 3 to 6 milligrams of caffeine per kilogram of body weight, or about 200 to 400 milligrams (2 to 3 cups of coffee) for a 150 lb. (68 kg) athlete about 30 minutes to an hour before exercise.

FUEL UTILIZATION

CAFFEINE METABOLISM

You will likely find that caffeine is particularly useful for endurance-based activity. Recently, British researchers found that relatively low doses of the stimulant, about two cups of coffee, could improve 40K time trial performance in trained cyclists by 55 to 84 seconds. In another study appearing in the *Journal of Pain*, exercise scientists found that cyclists who took caffeine before riding 30 minutes on stationary bikes had significantly less muscle pain during their effort than those who pedaled caffeine-free.

Though the research is a bit less conclusive, caffeine also may help improve strength performance. One review of 27 studies found that caffeine improved leg muscle power by up to 7 percent, but didn't seem to impact smaller muscle groups. It also appears to improve muscle endurance, so you can perform more repetitions for a given weight.

It may also boost performance in sprint activities. In a study on runners, those drinking regular coffee were 4.2 seconds faster in a 1500-meter run than those who were given decaf. And in studies on team sports, caffeine was shown to improve the passing accuracy in rugby and soccer sprint time.

Watch the dosage. Being a normal caffeine metabolizer does not make you immune to the potential negative side effects from too much caffeine, such as jitteriness and GI distress. So it is important to use it prudently.

One cup (8 oz.) of coffee generally delivers 100 to 150 mg of caffeine. Going above 600 mg can have adverse effects and may be prohibited by your sport's governing body. For endurance events lasting longer than two to three hours, you also may benefit from taking another dose of about 50 to 150 mg in the form of a caffeinated gel or energy drink for a boost during the latter part of the event.

Despite its longstanding reputation as such, caffeine is not a diuretic, so won't dehydrate you. As with any ergogenic aid, always be sure to test your response to caffeine in training before trying it during an event.

after about 90 minutes and starting to drop off after about 3 to 4 hours.

Caffeine eventually gets broken down in the liver by enzymes (Cytochrome P450 1A2, or CYP1A2) that metabolize the chemical. Depending on your genetic makeup, you will be able to metabolize caffeine at a normal rate, or your rate may be significantly slower. One study of 9,876 individuals found that variants in several genes were associated with slow caffeine metabolism (which was also associated with lower coffee consumption, indicating that people generally self regulate).

Being a slow caffeine metabolizer means the caffeine stays in your system longer, which can have adverse effects such as increasing blood pressure and may increase the risk of heart attack. Slow metabolizers also do not enjoy the same level of ergogenic improvement as people who metabolize the drug normally.


Our analysis investigated which genotype for these genes was present in your DNA. Your rating of **NORMAL** or **SLOW** reflects whether your genotype included those that carried a risk of adverse side effects in response to caffeine use or whether you are likely to benefit from using caffeine as an ergogenic aid.

RECOVERY & INJURY RISK

INJURY RISK

WHAT YOUR GENES SAY ABOUT YOU:

Our analysis indicates that your genetic profile exhibits characteristics that give you a likelihood of having an **ABOVE AVERAGE** fracture risk and a below average bone mineral density. That is concerning because weak bones not only leave you vulnerable to stress fractures and breaks, but also can be crippling and life-threatening later in life. As someone who is genetically inclined to have below average bone density, it's important to practice bone-building nutrition, exercise, and lifestyle habits.



Your genetic profile indicates that you are inclined to have **ABOVE AVERAGE** fracture risk.

You can minimize the potentially detrimental effects of your genes by eating, exercising, and practicing lifestyle habits that contribute to strong, healthy bone density.



RELATED GENES / SNPs

SPTBN1, MEPE, SLC25A13, MBL2/DKK1, LRP5, C18orf19

The genes and their associated SNPs in this category have been shown to have significant associations with a person's bone mineral density and risk of fracture.

Strong bones are essential for good health and longevity as well as to support your active lifestyle and avoid injuries like stress fractures and breaks. Throughout your life, your body is constantly breaking down old bone (in a process called resorption) and laying down new bone. You reach peak bone density at about age 30, after which you may start breaking down bone faster than you make it. If this loss remains unchecked, you can end up with dangerously thin bones that increase your risk of fracture. Because of hormonal reasons and the fact that they have thinner bones to begin with, women are at particular risk for thinning bones and osteoporosis (a disease where bones are porous and

SUCCESS STRATEGIES

As an active person, you're already on the right path to building a strong skeleton. But it's essential that you eat a diet that supports your activity level and encourages bone development. Your lifestyle habits and even certain medications can contribute to bone loss. So it's important to practice bone-building behaviors and avoid those (like smoking, which you likely already avoid) that can diminish your bone density.

Get all the calcium and vitamin D you need. Low calcium intake puts you at risk for low bone mineral density, as your bones are made mostly from this essential mineral. Men 70 years old and younger and women 50 years old and younger need 1,000 milligrams a day; men 71 years old and older and women 51 years old and older need 1,200 milligrams a day of calcium. You can get adequate calcium from dairy products like yogurt and cheese, sardines and salmon with bones, dark leafy vegetables, and tofu. Vitamin D assists in calcium absorption and bone formation. You get vitamin D through exposure to the sun and you can get it from fortified dairy products, eggs (with the yolks), and saltwater fish. Men and

RECOVERY & INJURY RISK

INJURY RISK

women 50 years old and younger need 400 to 800 IUs a day; men and women 51 years old and older need 800 to 1,000 IUs a day. Almost 70 percent of Americans don't meet all their vitamin D needs. You can ensure you get enough by taking a supplement of 400 to 800 IUs.

Prioritize protein. Active adults need more protein, which not only helps you build and maintain skeletal muscle, but also helps build your skeleton, as well. It's especially important to take in enough protein to fuel your training recovery, which in turn helps you maintain bone mineral density and testosterone, and for women can prevent menstrual cycle dysfunction, which can lead to significant bone loss. Aim for 30 to 40 grams per meal to help maintain muscle and bone integrity.

[IF FEMALE] *See your doctor if you stop menstruating and are not menopausal.* Irregular periods are a symptom of hormonal disruption that can cause bone loss. Amenorrheic athletes have up to a 30 percent lower bone mineral density as normally menstruating athletes. If your periods become irregular or you stop menstruating, see your doctor.

Train your core. Your spine is one of the most vulnerable spots for bone loss. Do core training, such as push-ups, pull ups, and planks year round.

Add impact and high intensity exercise. Impact exercise like walking, running, racquet sports, and jumping drills like plyometrics put healthy stress on your bones and encourage bone development. If you primarily practice non-impact sports like cycling and swimming, incorporate some impact cross training into your schedule. You may also build bone with low-impact exercise by adding high intensity, which research suggests may put enough tension on bones to help promote development.

Make muscle. Strong muscles and strong bones go hand in hand. Your muscles put healthy tension on your bones and encourage bone formation. Strength training is essential, especially once you hit your 40s, when muscle mass may naturally decline. It's also important to strength train those body parts that you don't use in your typical activity. Bone density is "site specific", meaning that you build more bone in the places that have the most muscle mass and that you work the hardest. Research has found that strength training can improve bone density by about 1 percent a year, according to the National Osteoporosis Foundation.

prone to breaks), especially after menopause.

Twins and family research reveals that up to 85 percent of the variance in bone mineral density (BMD) is determined by genetics.

The largest meta-analysis of 17 genome-wide association studies found that certain genetic scores were highly associated with BMD and fracture risk. The only way to know for certain that you have healthy or low BMD is to have a bone density test, called a dual energy x-ray absorptiometry or DXA scan, which measures bone density in your hip and spine. Other screenings, such as the kinds that measure bone density in your lower arm wrist, finger, or heel, also can identify thinning bones.

Our analysis investigated which genotype was present in your DNA. Your rating of **NORMAL**, **BELOW AVERAGE**, or **WELL BELOW AVERAGE** reflects whether or not your genotype includes those that increase your risk for low bone mineral density and bone fracture.

RECOVERY & INJURY RISK

INJURY RISK

Be cautious about bone-robbing medications. Certain medications such as corticosteroids, aluminum containing antacids, antidepressants, and proton pump inhibitors can contribute to bone loss. Talk to your doctor if you need to be on them for any length of time.

Go easy on alcohol. Heavy drinking can compromise bone integrity and lead to bone loss. If you drink alcohol, keep your consumption moderate: one drink a day for women, two for men.

Consider getting a bone scan. If you're concerned about the status of your bone health, make an appointment to get a bone scan. It's quick and easy and will let you know what, if any, actions you should be taking.

LINKS TO RELATED STUDIES

INTRINSIC MOTIVATION TO EXERCISE

J Behav Med. 2014 Dec;37(6):1180-92. doi: 10.1007/s10865-014-9567-4. Epub 2014 May 8.

What keeps a body moving? The brain-derived neurotrophic factor val66met polymorphism and intrinsic motivation to exercise in humans.

<https://www.ncbi.nlm.nih.gov/pubmed/?term=24805993>

Caldwell Hooper AE, Bryan AD, Hagger MS.

ADDICTIVE BEHAVIOR / STIMULUS CONTROL

Transl Psychiatry. 2015 Dec 1;5:e686. doi: 10.1038/tp.2015.176.

The significant association of Taq1A genotypes in DRD2/ANKK1 with smoking cessation in a large-scale meta-analysis of Caucasian populations

<https://www.ncbi.nlm.nih.gov/pubmed/?term=26624925>

Ma Y, Wang M, Yuan W, Su K, Li MD

POWER / ENDURANCE POTENTIAL

J Sci Med Sport. 2018 Feb;21(2):213-220. doi: 10.1016/j.jsams.2017.06.012. Epub 2017 Jun 21.

Nine genetic polymorphisms associated with power athlete status - A Meta-Analysis.

<https://www.ncbi.nlm.nih.gov/pubmed/?term=28666769>

Weyerstraß J, Stewart K, Wesselius A, Zeegers M.

GRIP STRENGTH / MUSCULAR FITNESS

Article number: 16015 (2017) doi:10.1038/ncomms16015

Large-scale GWAS identifies multiple loci for hand grip strength providing biological insights into muscular fitness

<https://www.nature.com/articles/ncomms16015>

Sara M. Willems, Daniel J. Wright, Robert A. Scott

TESTOSTERONE LEVELS

PLoS Genet. 2011 Oct;7(10):e1002313. doi: 10.1371/journal.pgen.1002313. Epub 2011 Oct 6.

Genetic determinants of serum testosterone concentrations in men.

<https://www.ncbi.nlm.nih.gov/pubmed/?term=21998597>

Ohlsson C, Wallaschowski H, Lunetta KL, Stolk L, Perry JR, Koster A, Petersen AK, Eriksson J, Lehtimäki T, Huhtaniemi IT, Hammond GL, Maggio M, Coviello AD; EMAS Study Group, Ferrucci L, Heier M, Hofman A, Holliday KL, Jansson JO, Kähönen M, Karasik D, Karlsson MK, Kiel DP, Liu Y, Ljunggren O, Lorentzon M, Lyytikäinen LP, Meitinger T, Mellström D, Melzer D, Miljkovic I, Nauck M, Nilsson M, Penninx B, Pye SR, Vasan RS, Reincke M, Rivadeneira F, Tajar A, Teumer A, Uitterlinden AG, Ulloor J, Viikari J, Völker U, Völzke H, Wichmann HE, Wu TS, Zhuang WV, Ziv E, Wu FC, Raitakari O, Eriksson A, Bidlingmaier M, Harris TB, Murray A, de Jong FH, Murabito JM, Bhasin S, Vandenput L, Haring R.

LINKS TO RELATED STUDIES

VO2 MAX

Physiol Genomics. 2003 Jul 7;14(2):161-6.

Associations between cardiorespiratory responses to exercise and the C34T AMPD1 gene polymorphism in the HERITAGE Family Study.

<https://www.ncbi.nlm.nih.gov/pubmed/12783984>

Rico-Sanz J, Rankinen T, Joannis DR, Leon AS, Skinner JS, Wilmore JH, Rao DC, Bouchard C; HERITAGE Family study.

Metabolism. 2004 Feb;53(2):193-202.

Apolipoprotein E genotype and changes in serum lipids and maximal oxygen uptake with exercise training.

<https://www.ncbi.nlm.nih.gov/pubmed/14767871>

Thompson PD, Tsongalis GJ, Seip RL, Bilbie C, Miles M, Zoeller R, Visich P, Gordon P, Angelopoulos TJ, Pescatello L, Bausserman L, Moyna N.

Metabolism. 2004 Jan;53(1):108-16.

Association of apolipoprotein E polymorphism with blood lipids and maximal oxygen uptake in the sedentary state and after exercise training in the HERITAGE family study.

<https://www.ncbi.nlm.nih.gov/pubmed/14681851>

Leon AS, Togashi K, Rankinen T, Després JP, Rao DC, Skinner JS, Wilmore JH, Bouchard C.

EXERCISE HEART RATE RESPONSE

Circ Cardiovasc Genet. 2010 Jun;3(3):294-9. doi: 10.1161/CIRCGENETICS.109.925644. Epub 2010 Apr 20.

CREB1 is a strong genetic predictor of the variation in exercise heart rate response to regular exercise: the HERITAGE Family Study.

<https://www.ncbi.nlm.nih.gov/pubmed/20407090>

Rankinen T1, Argyropoulos G, Rice T, Rao DC, Bouchard C.

EXERCISE STROKE VOLUME RESPONSE

Physiol Genomics. 2009 Jan 8;36(2):79-88. doi: 10.1152/physiolgenomics.00003.2008. Epub 2008 Nov 4.

KIF5B gene sequence variation and response of cardiac stroke volume to regular exercise.

<https://www.ncbi.nlm.nih.gov/pubmed/18984674>

Argyropoulos G, Stütz AM, Ilnytska O, Rice T, Teran-Garcia M, Rao DC, Bouchard C, Rankinen T.

BODY COMPOSITION RESPONSE TO STRENGTH TRAINING

International Journal of Obesity volume 39, pages 1371–1375 (2015) doi:10.1038/ijo.2015.78.

High genetic risk individuals benefit less from resistance exercise intervention

<http://www.nature.com/ijo/journal/vaop/ncurrent/abs/ijo201578a.html>

Y C Klimentidis, J W Bea, T Lohman, P-S Hsieh, S Going & Z Chen

LINKS TO RELATED STUDIES

PROTEIN UTILIZATION

Int J Obes (Lond). 2013 Dec;37(12):1545-52. doi: 10.1038/ijo.2013.54. Epub 2013 Apr 3.

FTO predicts weight regain in the Look AHEAD clinical trial.

<https://www.ncbi.nlm.nih.gov/pubmed/?term=23628854>

McCaffery JM, Papandonatos GD, Huggins GS, Peter I, Kahn SE, Knowler WC, Hudnall GE, Lipkin EW, Kitabchi AE, Wagenknecht LE, Wing RR; Genetic Subgroup of Look AHEAD; Look AHEAD Research Group.

FAT UTILIZATION

Diabetes Care. 2012 Feb;35(2):363-6. doi: 10.2337/dc11-1328. Epub 2011 Dec 16.

Genetic predictors of weight loss and weight regain after intensive lifestyle modification, metformin treatment, or standard care in the Diabetes Prevention Program.

<https://www.ncbi.nlm.nih.gov/pubmed/?term=22179955>

Delahanty LM, Pan Q, Jablonski KA, Watson KE, McCaffery JM, Shuldiner A, Kahn SE, Knowler WC, Florez JC, Franks PW; Diabetes Prevention Program Research Group.

Diabetes. 2002 Aug;51(8):2581-6.

Association of the Pro12Ala polymorphism in the PPAR-gamma2 gene with 3-year incidence of type 2 diabetes and body weight change in the Finnish Diabetes Prevention Study.

<https://www.ncbi.nlm.nih.gov/pubmed/?term=12145174>

Lindi VI, Uusitupa MI, Lindström J, Louheranta A, Eriksson JG, Valle TT, Hämäläinen H, Ilanne-Parikka P, Keinänen-Kiukaanniemi S, Laakso M, Tuomilehto J; Finnish Diabetes Prevention Study.

Clin Genet. 2003 Feb;63(2):109-16.

The PPAR-gamma P12A polymorphism modulates the relationship between dietary fat intake and components of the metabolic syndrome: results from the Québec Family Study.

<https://www.ncbi.nlm.nih.gov/pubmed/?term=12630956>

Robitaille J, Després JP, Périusse L, Vohl MC.

Hum Mol Genet. 2003 Nov 15;12(22):2923-9. Epub 2003 Sep 23.

Interaction between a peroxisome proliferator-activated receptor gamma gene polymorphism and dietary fat intake in relation to body mass.

<https://www.ncbi.nlm.nih.gov/pubmed/?term=14506127>

Memisoglu A, Hu FB, Hankinson SE, Manson JE, De Vivo I, Willett WC, Hunter DJ.

Am J Clin Nutr. 2012 Nov;96(5):1129-36. doi: 10.3945/ajcn.112.038125. Epub 2012 Oct 3.

TCF7L2 genetic variants modulate the effect of dietary fat intake on changes in body composition during a weight-loss intervention.

<https://www.ncbi.nlm.nih.gov/pubmed/?term=23034957>

Mattei J, Qi Q, Hu FB, Sacks FM, Qi L.

Circulation. 2006 May 2;113(17):2062-70. Epub 2006 Apr 24.

Dietary intake of n-6 fatty acids modulates effect of apolipoprotein A5 gene on plasma fasting triglycerides, remnant lipoprotein concentrations, and lipoprotein particle size: the Framingham Heart Study.

<https://www.ncbi.nlm.nih.gov/pubmed/?term=16636175>

Lai CQ, Corella D, Demissie S, Cupples LA, Adiconis X, Zhu Y, Parnell LD, Tucker KL, Ordovas JM.

LINKS TO RELATED STUDIES

Clin Genet. 2005 Aug;68(2):152-4.

A polymorphism in the apolipoprotein A5 gene is associated with weight loss after short-term diet.

<https://www.ncbi.nlm.nih.gov/pubmed/?term=15996212>

Aberle J, Evans D, Beil FU, Seedorf U.

J Mol Med (Berl). 2007 Feb;85(2):119-28. Epub 2007 Jan 9.

APOA5 gene variation modulates the effects of dietary fat intake on body mass index and obesity risk in the Framingham Heart Study.

<https://www.ncbi.nlm.nih.gov/pubmed/?term=17211608>

Corella D, Lai CQ, Demissie S, Cupples LA, Manning AK, Tucker KL, Ordovas JM.

J Nutr. 2011 Mar;141(3):380-5. doi: 10.3945/jn.110.130344. Epub 2011 Jan 5.

APOA5 gene variation interacts with dietary fat intake to modulate obesity and circulating triglycerides in a Mediterranean population.

<https://www.ncbi.nlm.nih.gov/pubmed/?term=21209257>

Sánchez-Moreno C, Ordovás JM, Smith CE, Baraza JC, Lee YC, Garaulet M.

Am J Clin Nutr. 2014 Feb;99(2):392-9. doi: 10.3945/ajcn.113.072066. Epub 2013 Dec 11.

Variants in glucose- and circadian rhythm-related genes affect the response of energy expenditure to weight-loss diets: the POUNDS LOST Trial.

<https://www.ncbi.nlm.nih.gov/pubmed/?term=24335056>

Mirzaei K, Xu M, Qi Q, de Jonge L, Bray GA, Sacks F, Qi L.

Circulation. 2013 Mar 26;127(12):1283-9. doi: 10.1161/CIRCULATIONAHA.112.000586. Epub 2013 Feb 27.

Genetic determinant for amino acid metabolites and changes in body weight and insulin resistance in response to weight-loss diets: the Preventing Overweight Using Novel Dietary Strategies (POUNDS LOST) trial.

<https://www.ncbi.nlm.nih.gov/pubmed/?term=23446828>

Xu M, Qi Q, Liang J, Bray GA, Hu FB, Sacks FM, Qi L.

CARB UTILIZATION

Circulation. 2011 Aug 2;124(5):563-71. doi: 10.1161/CIRCULATIONAHA.111.025767. Epub 2011 Jul 11.

Insulin receptor substrate 1 gene variation modifies insulin resistance response to weight-loss diets in a 2-year randomized trial: the Preventing Overweight Using Novel Dietary Strategies (POUNDS LOST) trial.

<https://www.ncbi.nlm.nih.gov/pubmed/?term=21747052>

Qi Q, Bray GA, Smith SR, Hu FB, Sacks FM, Qi L.

CAFFEINE METABOLISM

Hum Mol Genet. 2016 Dec 15;25(24):5472-5482. doi: 10.1093/hmg/ddw334.

Genome-wide association study of caffeine metabolites provides new insights to caffeine metabolism and dietary caffeine-consumption behavior.

<https://www.ncbi.nlm.nih.gov/pubmed/?term=27702941>

Cornelis MC1, Kacprowski T2,3, Menni C4, Gustafsson S5, Pivn E6, Adamski J7,8,9, Artati A7, Eap CB10,11, Ehret G12,13, Friedrich N3,14, Ganna A15,16, Guessous I6,17,18, Homuth G2, Lind L19, Magnusson PK20, Mangino M4, Pedersen NL20, Pietzner M3,14, Suhre K21,22, Völzke H23; Swiss Kidney Project on Genes in Hypertension (SKIPOGH) team, Bochud M6, Spector TD4, Grabe HJ24, Ingelsson E5,25.

LINKS TO RELATED STUDIES

SYSTEMIC INFLAMMATION

Circulation. 2011 Feb 22;123(7):731-8. doi: 10.1161/CIRCULATIONAHA.110.948570. Epub 2011 Feb 7.

Meta-analysis of genome-wide association studies in >80 000 subjects identifies multiple loci for C-reactive protein levels.

<https://www.ncbi.nlm.nih.gov/pubmed/?term=21300955>

Dehghan A, Dupuis J, Barbalic M, Bis JC, Eiriksdottir G, Lu C, Pellikka N, Wallaschofski H, Kettunen J, Henneman P, Baumert J, Strachan DP, Fuchsberger C, Vitart V, Wilson JF, Paré G, Naitza S, Rudock ME, Surakka I, de Geus EJ, Alizadeh BZ, Guralnik J, Shuldiner A, Tanaka T, Zee RY, Schnabel RB, Nammi V, Kavousi M, Ripatti S, Nauck M, Smith NL, Smith AV, Sundvall J, Scheet P, Liu Y, Ruokonen A, Rose LM, Larson MG, Hoogeveen RC, Freimer NB, Teumer A, Tracy RP, Launer LJ, Buring JE, Yamamoto JF, Folsom AR, Sijbrands EJ, Pankow J, Elliott P, Keaney JF, Sun W, Sarin AP, Fontes JD, Badola S, Astor BC, Hofman A, Pouta A, Werdan K, Greiser KH, Kuss O, Meyer zu Schwabedissen HE, Thiery J, Jamshidi Y, Nolte IM, Soranzo N, Spector TD, Völzke H, Parker AN, Aspelund T, Bates D, Young L, Tsui K, Siscovick DS, Guo X, Rotter JI, Uda M, Schlessinger D, Rudan I, Hicks AA, Penninx BW, Thorand B, Gieger C, Coresh J, Willemsen G, Harris TB, Uitterlinden AG, Järvelin MR, Rice K, Radke D, Salomaa V, Willems van Dijk K, Boerwinkle E, Vasani RS, Ferrucci L, Gibson QD, Bandinelli S, Snieder H, Boomsma DI, Xiao X, Campbell H, Hayward C, Pramstaller PP, van Duijn CM, Peltonen L, Psaty BM, Gudnason V, Ridker PM, Homuth G, Koenig W, Ballantyne CM, Witteman JC, Benjamin EJ, Perola M, Chasman DI.

INJURY RISK

Nat Genet. 2012 Apr 15;44(5):491-501. doi: 10.1038/ng.2249.

Genome-wide meta-analysis identifies 56 bone mineral density loci and reveals 14 loci associated with risk of fracture.

<https://www.ncbi.nlm.nih.gov/pubmed/?term=22504420>

Estrada K, Styrkarsdottir U, Evangelou E, Hsu YH, Duncan EL, Ntzani EE, Oei L, Albagha OM, Amin N, Kemp JP, Koller DL, Li G, Liu CT, Minster RL, Moayyeri A, Vandenput L, Willner D, Xiao SM, Yerges-Armstrong LM, Zheng HF, Alonso N, Eriksson J, Kammerer CM, Kaptoge SK, Leo PJ, Thorleifsson G, Wilson SG, Wilson JF, Aalto V, Alen M, Aragaki AK, Aspelund T, Center JR, Dailiana Z, Duggan DJ, Garcia M, Garcia-Giralt N, Giroux S, Hallmans G, Hocking LJ, Husted LB, Jameson KA, Khusainova R, Kim GS, Kooperberg C, Koromila T, Kruk M, Laaksonen M, Lacroix AZ, Lee SH, Leung PC, Lewis JR, Masi L, Mencej-Bedrac S, Nguyen TV, Nogues X, Patel MS, Prezelj J, Rose LM, Scollen S, Siggeirsdottir K, Smith AV, Svensson O, Trompet S, Trummer O, van Schoor NM, Woo J, Zhu K, Balcells S, Brandi ML, Buckley BM, Cheng S, Christiansen C, Cooper C, Dedoussis G, Ford I, Frost M, Goltzman D, González-Macías J, Kähönen M, Karlsson M, Khusnutdinova E, Koh JM, Kollia P, Langdahl BL, Leslie WD, Lips P, Ljunggren Ö, Lorenc RS, Marc J, Mellström D, Obermayer-Pietsch B, Olmos JM, Pettersson-Kymmer U, Reid DM, Riancho JA, Ridker PM, Rousseau F, Slagboom PE, Tang NL, Urreizti R, Van Hul W, Viikari J, Zarrabeitia MT, Aulchenko YS, Castano-Betancourt M, Grundberg E, Herrera L, Ingvarsson T, Johannsdottir H, Kwan T, Li R, Luben R, Medina-Gómez C, Palsson ST, Reppe S, Rotter JI, Sigurdsson G, van Meurs JB, Verlaan D, Williams FM, Wood AR, Zhou Y, Gautvik KM, Pastinen T, Raychaudhuri S, Cauley JA, Chasman DI, Clark GR, Cummings SR, Danoy P, Dennison EM, Eastell R, Eisman JA, Gudnason V, Hofman A, Jackson RD, Jones G, Jukema JW, Khaw KT, Lehtimäki T, Liu Y, Lorentzon M, McCloskey E, Mitchell BD, Nandakumar K, Nicholson GC, Oostra BA, Peacock M, Pols HA, Prince RL, Raitakari O, Reid IR, Robbins J, Sambrook PN, Sham PC, Shuldiner AR, Tylavsky FA, van Duijn CM, Wareham NJ, Cupples LA, Econs MJ, Evans DM, Harris TB, Kung AW, Psaty BM, Reeve J, Spector TD, Streeten EA, Zillikens MC, Thorsteinsdottir U, Ohlsson C, Karasik D, Richards JB, Brown MA, Stefansson K, Uitterlinden AG, Ralston SH, Ioannidis JP, Kiel DP, Rivadeneira F.